

To: (10)(2e) [(10)(2e)]@who.int
Cc: (10)(2e) [(10)(2e)]@who.int; (10)(2e) [(10)(2e)]@who.int
From: (10)(2e); (10)(2e)
Sent: Fri 9/4/2020 2:25:50 PM
Subject: RE: query on quantitative serology assay
Received: Fri 9/4/2020 2:25:51 PM
[20200902 Tables final.pdf](#)
[20200902 Figures ELISA NT final.pdf](#)
[20200902 Genescript NT paper final.pdf](#)

Hi,

What context? Various labs use various tests.
Dutch government has stockpile of Wantai tests , this is used most. Now more CLIA are on the market more labs are validating this.

At RIVM we use Wantai total Ab, VNT, multiplex micro-array, genescript pseudoVNT. Please find attached a manuscript we just submitted together with (10)(2e) (10)(2e)

Cheers (10)(2e)

From: (10)(2e) <(10)(2e)]@who.int>
Sent: vrijdag 4 september 2020 14:36
To: (10)(2e); (10)(2e) <(10)(2e)]@rivm.nl>
Cc: (10)(2e) <(10)(2e)]@who.int>; (10)(2e) <(10)(2e)]@who.int>
Subject: query on quantitative serology assay

Hi (10)(2e)

I was wondering: which test do you use to look at Ab titers? Are you relying only on PRNT or you have a quantitative ELISA/CLIA that you use for binding antibody titers?

Thanks in advance,

(10)(2e)