

To: [redacted] [redacted] [redacted]@rivm.nl]
Cc: [redacted] [redacted] [redacted]@sciensano.be]; [redacted] [redacted] [redacted]@sciensano.be]
From: [redacted] [redacted] [redacted]@sciensano.be]
Sent: Wed 9/23/2020 12:30:05 PM
Subject: RE: oracol swabs en SARS-CovID
Received: Wed 9/23/2020 12:30:18 PM

Hi [redacted]

Thank you very much for your quick reply.

Very interesting findings and happy to discuss this further with [redacted]

For PCR testing, you perform two Oracol samples in order to obtain sufficient saliva (minimal 2ml).
We tested it by ourselves and indeed it is not easy to collect 1 ml with only 1 Oracol device.

Probably I will come back to you with some extra questions.

Kind regards

[redacted]

[redacted] [redacted], [redacted]
[redacted]

Viral Diseases

National Reference Center for Measles, Mumps, Rubella, Viral Hepatitis and Human Papillomavirus

T + 32 [redacted]



Rue Juliette Wytsmanstraat 14 • 1050 Brussels
Laboratory adress : Rue Engelandstraat 642 • 1180 Brussels

www.sciensano.be

Please consider the environment before printing this e-mail

[redacted]

[redacted]

[redacted]

[redacted]

[redacted]

[redacted]

[redacted]

[redacted]

Dubbel

Dubbel

5.1.2e 5.1.2e

5.1.2e 5.1.2e

5.1.2e 5.1.2e

5.1.2e

5.1.2e

5.1.2e

5.1.2e

Dubbel

5.1.2e

5.1.2e

Dubbel