To: 12. 5.1.2e [5.1.2e @minvws.nl]

From: H.M. de Jonge

Sent: Wed 2/3/2021 7:55:33 PM

Subject: Fwd: Goed nieuws over langere tijd tussen prime-boost + reductie van transmissie

Received: Wed 2/3/2021 7:55:50 PM

Verstuurd vanaf mijn iPad

Begin doorgestuurd bericht:

Van: 5.1.2e < 5.1.2e @radboudumc.nl>

Datum: 3 februari 2021 om 13:10:49 CET

Aan: 5.1.2e @icloud.com

Onderwerp: Goed nieuws over langere tijd tussen prime-boost + reductie van transmissie

Ha Huug, goed nieuws; met uitstellen van de booster vaccinatie gaat effectiviteit van bescheming juist omhoog, daarnaast is voor het eerst aangetoond dat gebruik van AstraZeneca-Oxford vaccin leidt tot reductie van transmissie. Nu maar hopen dat ze op tijd kunnen leveren .

Groeten,

5.1.2e

```
From: 5.1.2e 5.1.2e < 5.1.2e @phe.gov.uk>
Sent: dinsdag 2 februari 2021 19:08
To: 5.1.2e
                           @bristol.ac.uk>; 5.1.2e @ftcon.co.uk;
                                                                                @wales.nhs.uk; 5.1.2e
         5.1.2e
                                                       5.1.2e
                                                                @lshtm.ac.uk';
                                 @phc.ox.ac.uk>;
              5.1.2e
                                                          <
                        @dhsc.gov.uk>; 5.1.2e
                                                                            @pirbright.ac.uk
                        @rivm.nl';
                                                          5.1.2e
                                                                     @phe.gov.uk>;
                     @ucl.ac.uk>; 5.1.2e 5.1.2e
                                                                  @hscni.net>;
                                                          @nhs.net>; 5.1.2e 5.1.2e
                   @phe.gov.uk>;
                                             <
                     <u>@phe.gov.uk</u>>; 5.1.2e , 12 5.1.2e < 5.1.2e <u>@imperial.ac.uk</u>>;
                                          5.1.2e
                                                    @nhslothian.scot.nhs.uk; 5.1.2e 5.1.2e
                      @nuh.nhs.uk>;
                                               @warwick.ac.uk';
                    @bristol.ac.uk>;
                                                                            @rug.nl;
                                                                                                   @nhs.net';
                                  5.1.2e @soton.ac.uk'; 5.1.2e
                   @gmail.com;
                                                                                            @phe.gov.uk>;
                                       @ntlworld.com>; 5.1.2e
                              5.1.2e
                          @UHBristol.nhs.uk>
Cc: 5.1.2e 5.1.2e <
                                                                             @phe.gov.uk>; 5.1.2e 5.1.2e
                             @phe.gov.uk>; 5.1.2e 5.1.2e
                       @phe.gov.uk>
```

Subject: [Spam] Oxford coronavirus vaccine shows sustained protection of 76% during the 3-month interval until the second dose

FOR IMMEDIATE RELEASE

University of Oxford Press Release

Oxford coronavirus vaccine shows sustained protection of 76% during the 3-month interval until the second dose

 Analyses reveal single standard dose efficacy from day 22 to day 90 post vaccination of 76% with protection not falling in this three-month period

- After the second dose vaccine efficacy from two standard doses is 82.4% with the 3-month interval being used in the UK. (82.4% effective, with a 95% confidence interval of 62.7% 91.7% at 12+ weeks)
- Data supports the 4-12 week prime-boost dosing interval recommended by many global regulators
- Analyses of PCR positive swabs in UK population suggests vaccine may have substantial effect on transmission of the virus with 67% reduction in positive swabs among those vaccinated

Researchers at the University of Oxford have today published in <u>Preprints with The Lancet</u> an analysis of further data from the ongoing trials of the vaccine. In this, they reveal that the vaccine efficacy is higher at longer prime-boost intervals, and that a single dose of the vaccine is 76% effective from 22- to up to 90-days post vaccination.

In this preprint, which is currently under review at The Lancet, they report on an analysis of additional data to include information from the trial up to the 7th December 2020, which includes a further 201 cases of primary symptomatic COVID-19 (332 cases from 131 reported in previously), They report that the effect of dosing interval on efficacy is pronounced, with <u>vaccine efficacy rising from 54.9% with an interval of less than six weeks to 82.4% when spaced 12 or more weeks apart.</u>

They also detail that a single standard dose of the vaccine is 76% effective at protecting from primary symptomatic COVID-19 for the first 90 days post vaccination, once the immune system has built this protection 22 days after the vaccination, with the protection showing little evidence of waning in this period.

Professor 5.1.2e Chief Investigator of the Oxford Vaccine Trial, and co-author of the paper, said:

'These new data provide an important verification of the interim data that was used by more than 25 regulators including the MHRA and EMA to grant the vaccine emergency use authorisation.

'It also supports the policy recommendation made by the Joint Committee on Vaccination and Immunisation (JCVI) for a 12-week prime-boost interval, as they look for the optimal approach to roll out, and reassures us that people are protected from 22 days after a single dose of the vaccine.'

The exploratory analyses presented in this preprint paper suggest that it is the dosing interval and not the dosing level which has a great impact on the efficacy of the vaccine. This is in line with previous research supporting greater efficacy with longer prime-boost intervals done with other vaccines such as influenza, Ebola and malaria.

The authors also report further on the potential for the vaccine to reduce transmission of the virus, based on swabs obtained from volunteers in the UK arms of the trial with **a 67% reduction** after the first dose of the vaccine.

They also hope to report data regarding the new variants in the coming days, and expect the findings to be broadly similar to those already reported by fellow vaccine developers.

ENDS

Notes to editors:

For further information or to arrange an interview, please contact the University of Oxford press office at 6.1.2e @admin.ox.ac.uk or on 5.1.2e

For more about the Oxford vaccine project and team: www.ox.ac.uk/covid-vaccine

De informatie in dit bericht is uitsluitend bestemd voor de geadresseerde. Aan dit bericht en de bijlagen kunnen geen rechten worden ontleend. Heeft u deze e-mail onbedoeld ontvangen? Dan verzoeken wij u het te vernietigen en de afzender te informeren. Openbaar maken, kopiëren en verspreiden van deze e-mail of informatie uit deze e-mail is alleen toegestaan met voorafgaande schriftelijke toestemming van de afzender. Het Radboudumc staat geregistreerd bij de Kamer van Koophandel in het handelsregister onder nummer 80262783.

The content of this message is intended solely for the addressee. No rights can be derived from this message or its attachments. If you are not the intended recipient, we kindly request you to delete the message and inform the sender. It is strictly prohibited to disclose, copy or distribute this email or the information inside it, without a written consent from the sender. Radboud university medical center is registered with the Dutch Chamber of Commerce trade register with number 80262783.