

To: [redacted] 5.1.2e [redacted] 5.1.2e @rivm.nl]
From: [redacted] 5.1.2e
Sent: Thur 4/30/2020 5:38:25 PM
Subject: RE: guidelines
Received: Thur 4/30/2020 5:38:25 PM

Dank dit neem ik zeker mee naar [redacted] 5.1.2e . Uit welk van de doc heb jij dit gehaald van [redacted] 5.1.2e ?

Hartelijke groet,

[redacted] 5.1.2e

Van: [redacted] 5.1.2e <[redacted] 5.1.2e @rivm.nl>
Verzonden: donderdag 30 april 2020 16:35
Aan: [redacted] 5.1.2e <[redacted] 5.1.2e @rivm.nl>
Onderwerp: guidelines

Hoi [redacted] 5.1.2e ,

In een van de documenten die [redacted] 5.1.2e stuurde stond volgende, kunnen we wel wat mee, vooral bij checken of LCI zo compleet mogelijk is, toch?

4. Guidance for Vaccination Posts

- Conduct vaccination sessions in well-ventilated areas that are frequently disinfected. (13)
- Ensure the availability of hand sanitizer or a hand washing station with chlorinated water for use by users at the entrance of the health facility. (13)
- Limit the number of family members accompanying the person to be vaccinated (one companion).
- Perform triage of persons presenting respiratory symptoms before admission to the vaccination posts to prevent the spread of SARS-CoV-2. If patient presents respiratory symptoms, offer medical mask, do not vaccinate, and refer to service for evaluation.
- Avoid crowded waiting rooms. Some strategies for this could include:
 - Scheduled times for vaccination appointments;
 - Integrate vaccination activities with other essential preventive health services, as appropriate;
 - Carry out small and frequent vaccination sessions;
 - Utilize outdoor spaces and adhere to the recommendation of social distance within the facility, or vaccination post;
 - Establish exclusive vaccination sessions for older people and people with pre-existing medical conditions (such as high blood pressure, heart disease, respiratory disease, or diabetes).
- Whenever possible, the vaccination post must be separate from healing services (i.e., different hours, different spaces);
- Recommendations for vaccinators:
 - Frequently maintain hand hygiene as described in "My 5 Moments for Hand Hygiene": i) Before touching a patient; ii) Before clean or aseptic procedures; iii) After body fluid exposure/risk; iv) After touching a patient, and v) After touching patient surroundings. (14)
 - Hand hygiene consists of washing hands with soap and water or with a hand sanitizer that contains between 60% and 80% alcohol.
 - Comply with guidelines on clothing: i) Use a uniform, which should be not used outside the health facility; ii) Use closed shoes; iii) Do not use accessories (for example: earrings, rings, chains, watch).
 - Clean cell phones properly. Do not use cell phone while providing medical attention.
 - If you experience symptoms, such as cough or fever, you should not be working and should seek medical attention.

[redacted] 5.1.2e

[redacted] 5.1.2e

RIVM Centrum voor Bevolkingsonderzoek | Postbus 1 | 3720 BA Bilthoven

M: [redacted] 5.1.2e | **E:** [redacted] 5.1.2e @rivm.nl | [redacted] 5.1.2e

[redacted] 5.1.2e