

To: 5.1.2e [5.1.2e @rivm.nl]
Cc: 5.1.2e [5.1.2e @rivm.nl]; 5.1.2e [5.1.2e @rivm.nl]
From: 5.1.2e
Sent: Tue 5/19/2020 11:00:39 AM
Subject: FW: First results of the RCGP study in England
Received: Tue 5/19/2020 11:00:40 AM

FYI: an article from our UK colleagues, in which they used a similar method as in the I-MOVE Covid19 Risk factor protocol.
Probably this article is a bit more to the point than the huge I-MOVE protocols.

I realize that the I-MOVE protocols (as well as this article) are very specific for the primary care setting, as 5.1.2e and 5.1.2e already indicated. So probably it is not a perfect parallel. Maybe we can pick some ideas for interesting variables from the list of variables in the appendix of the protocol, but for the rest I think it might be better to develop a totally new protocol from scratch (with some of the ideas from 5.1.2e) for the setting of the testing streets. That's how I see it now. We can discuss about this later this afternoon.

Best regards,

5.1.2e

5.1.2e

5.1.2e

5.1.2e

5.1.2e

5.1.2e

5.1.2e

5.1.2e

5.1.2e

5.1.2e

5.1.2e

5.1.2e

5.1.2e

5.1.2e

5.1.2e

5.1.2e

5.1.2e

Dubbel

5.1.2e

2 - 2

Dubbel